

New Higher-Derivative R^4 Theorems

Nathan Berkovits¹

*Instituto de Física Teórica, State University of São Paulo
Rua Pamplona 145, 01405-900, São Paulo, SP, Brasil*

The non-minimal pure spinor formalism for the superstring is used to prove two new multiloop theorems which are related to recent higher-derivative R^4 conjectures of Green, Russo and Vanhove. The first theorem states that when $0 < n < 12$, $\partial^n R^4$ terms in the Type II effective action do not receive perturbative contributions above $n/2$ loops. The second theorem states that when $n \leq 8$, perturbative contributions to $\partial^n R^4$ terms in the IIA and IIB effective actions coincide.

August 2006

¹ e-mail: nberkovi@ift.unesp.br

1. Introduction

Computation of superstring multiloop amplitudes is useful for verifying perturbative finiteness and for testing non-perturbative duality conjectures. Although there are several prescriptions available for computing multiloop amplitudes, the most efficient prescription is based on the pure spinor formalism which is manifestly super-Poincaré covariant and which involves a bosonic ghost λ^α satisfying the pure spinor constraint $\lambda\gamma^m\lambda = 0$ [1].

Using the minimal version of the pure spinor formalism, a multiloop prescription was defined in [2] and used to prove certain vanishing theorems. One theorem stated that massless N -point multiloop amplitudes are vanishing when $N \leq 3$, which is related to perturbative finiteness of the superstring [3]. Another theorem stated that R^4 terms in the Type II effective action do not receive perturbative contributions above one-loop, which is related to S-duality of the Type IIB superstring [4][5]. In later papers, four-point one and two-loop amplitudes were computed using the minimal formalism and were shown to coincide with the RNS result [6].

Recently, a new multiloop prescription was proposed using a non-minimal version of the pure spinor formalism which involves both the pure spinor λ^α and its complex conjugate $\bar{\lambda}_\alpha$ [7]. This non-minimal prescription for multiloop amplitudes has several advantages over the minimal prescription. Firstly, unlike in the minimal formalism, the non-minimal formalism allows the construction of a composite b ghost satisfying $\{Q, b\} = T$. Secondly, the non-minimal formalism can be interpreted as a critical topological string, so the non-minimal amplitude prescription is the same as in bosonic string theory. Thirdly, there is no need for picture-changing operators in the non-minimal prescription, which were inconvenient in the minimal prescription since they broke manifest Lorentz covariance at intermediate stages in the computation.

The only difficulty in the non-minimal prescription is regularizing the functional integral over the pure spinor ghosts when $\lambda \rightarrow 0$, which will be discussed in an upcoming paper with Nikita Nekrasov [8]. Fortunately, this $\lambda \rightarrow 0$ regularization is unnecessary for proving the multiloop theorems in this paper. Furthermore, it was recently shown with Carlos Mafra that various one and two-loop amplitude computations using the non-minimal prescription correctly reproduce the RNS result [9]. Although it should be possible to directly prove the equivalence of the minimal and non-minimal amplitude prescriptions (perhaps using the Cech description of Nekrasov [10]), this has not yet been done.

In this paper, the non-minimal prescription will be used to prove two new multiloop theorems which are related to recent higher-derivative R^4 conjectures of Green, Russo and

Vanhove based on duality symmetries [11] [12] [13] [14] [15] [16]. The first new multiloop theorem states that when $0 < n < 12$, $\partial^n R^4$ terms in the Type II effective action do not receive genus g contributions for $g > n/2$. The restriction that $n < 12$ is related to the fact that $\partial^{12} R^4$ can be written as a superspace integral over 32 θ 's. The second new multiloop theorem states that when $n \leq 8$, perturbative contributions to $\partial^n R^4$ terms in the IIA and IIB effective action coincide. For $n \leq 4$, this can be shown using the RNS formalism [14], and for $n = 6$, it was recently conjectured by Green and Vanhove [12].

In section 2, the amplitude prescription using the non-minimal formalism will be reviewed and, in section 3, the two new multiloop theorems will be proven.

2. Review of Non-Minimal Amplitude Prescription

2.1. Amplitude prescription

Using the non-minimal pure spinor formalism, the N -point g -loop amplitude prescription is [7]

$$\mathcal{A} = \int d^{6g-6} \tau \langle | \prod_{j=1}^{3g-3} (\int dy_j \mu_j(y_j) b(y_j)) |^2 \prod_{r=1}^N \int d^2 z_r U_r(z_r) |\mathcal{N}|^2 \rangle \quad (2.1)$$

where $\int d^{6g-6} \tau \langle | \prod_{j=1}^{3g-3} (\int dy_j \mu_j(y_j) b(y_j)) |^2 \prod_{r=1}^N \int d^2 z_r U_r(z_r) \rangle$ is the usual amplitude prescription of bosonic string theory, τ_j and μ_j are the Teichmüller parameters and corresponding Beltrami differentials, b is a composite operator satisfying $\{Q, b\} = T$, and $\mathcal{N} = e^{\{Q, \chi\}}$ is a BRST-invariant operator which regularizes the 0/0 coming from integration over the worldsheet zero modes.

The left-moving worldsheet variables in the non-minimal pure spinor formalism include the Green-Schwarz-Siegel variables $(x^m, \theta^\alpha, d_\alpha)$, the pure spinor ghosts $(\lambda^\alpha, w_\alpha)$, and the non-minimal variables $(\bar{\lambda}_\alpha, \bar{w}^\alpha, r_\alpha, s^\alpha)$. In terms of these variables, the composite b operator is

$$b = s^\alpha \partial \bar{\lambda}_\alpha + \frac{\bar{\lambda}_\alpha (2\Pi^m (\gamma_m d)^\alpha - N_{mn} (\gamma^{mn} \partial \theta)^\alpha - J \partial \theta^\alpha - \frac{1}{4} \partial^2 \theta^\alpha)}{4(\bar{\lambda} \lambda)} \quad (2.2)$$

$$+ \frac{(\bar{\lambda} \gamma^{mnp} r)(d \gamma_{mnp} d + 24 N_{mn} \Pi_p)}{192(\bar{\lambda} \lambda)^2} - \frac{(r \gamma_{mnp} r)(\bar{\lambda} \gamma^m d) N^{np}}{16(\bar{\lambda} \lambda)^3} + \frac{(r \gamma_{mnp} r)(\bar{\lambda} \gamma^{pqr} r) N^{mn} N_{qr}}{128(\bar{\lambda} \lambda)^4}$$

where $\Pi^m = \partial x^m + \theta \gamma^m \partial \theta$ is the supersymmetric momentum, and $N_{mn} = \frac{1}{2} w \gamma_{mn} \lambda$ and $J = \lambda w$ are the Lorentz and ghost currents for the pure spinors. Although the expression

of (2.2) is complicated, it will turn out that only two terms in (2.2) are relevant for proving the multiloop theorems, namely the terms $\frac{\Pi^m(\bar{\lambda}\gamma_m d_\alpha)}{2(\lambda\bar{\lambda})}$ and $\frac{(\bar{\lambda}\gamma^{mnp}r)(d\gamma_{mnp}d)}{192(\lambda\bar{\lambda})^2}$.

The zero-mode regulator $\mathcal{N}$ in (2.1) can be defined as

$$\mathcal{N} = \exp\{Q, \chi\} \quad (2.3)$$

$$= \exp[-\lambda^\alpha \bar{\lambda}_\alpha - r_\alpha \theta^\alpha - \frac{1}{2} N_{mn}^I \bar{N}^{mnI} - J^I \bar{J}^I - \frac{1}{4} (s^I \gamma^{mn} \bar{\lambda})(d^I \gamma_{mn} \lambda) - (\bar{\lambda} s^I)(\lambda d^I)],$$

where $s^{\alpha I} = \oint_{a_I} dz s^\alpha$, $d_\alpha^I = \oint_{a_I} dz d_\alpha$, $N_{mn}^I = \oint_{a_I} dz N_{mn}$, $J^I = \oint_{a_I} dz J$ are the zero modes obtained by integrating these fields around the I^{th} a -cycle, and

$$\chi = -\bar{\lambda}_\alpha \theta^\alpha - \frac{1}{2} N_{mn}^I (s^I \gamma^{mn} \bar{\lambda}) - J^I (\bar{\lambda} s^I). \quad (2.4)$$

When the bosonic zero modes of λ^α , N_{mn}^I or J^I go to infinity, the functional integral over these zero modes is well-defined because of the exponential cutoff in $\mathcal{N}$.

However, when $\lambda^\alpha \rightarrow 0$, the poles in (2.2) make the functional integral over λ^α ill-defined if the sum of the degree of the poles is greater than or equal to 11. If the contributions from the b ghosts diverge as fast as $(\lambda\bar{\lambda})^{-11}$, the measure factor $\int d^{11} \lambda d^{11} \bar{\lambda}$ does not converge fast enough to make the functional integral well-defined. In an upcoming paper with Nekrasov [8], it will be shown how to regularize this $\lambda \rightarrow 0$ divergence for arbitrary multiloop amplitudes. However, there are certain amplitudes for which the sum of the degree of the poles from the b ghosts is always less than 11, so one does not need to worry about regularizing the $\lambda \rightarrow 0$ divergence. For example, since the maximum pole in the b ghost is of degree 3, there is no $\lambda \rightarrow 0$ divergence when the genus is less than or equal to two since, for these amplitudes, there are three or fewer b ghosts. As will also be discussed in [8], another type of amplitude for which there is no $\lambda \rightarrow 0$ divergence is when at least one of the sixteen θ^α zero modes comes from $\mathcal{N}$.

2.2. F -terms

To show that terms receiving at least one θ zero mode from $\mathcal{N}$ do not contain divergences when $\lambda \rightarrow 0$, first note that BRST invariance implies that the $e^{-(\lambda\bar{\lambda}+r\theta)}$ term in $\mathcal{N}$ can be modified to $e^{-\rho(\lambda\bar{\lambda}+r\theta)}$ for any positive constant ρ . Because

$$e^{-\rho(\lambda\bar{\lambda}+r\theta)} = e^{\{Q, -\rho\theta\bar{\lambda}\}} = 1 + \{Q, \xi_\rho\} \quad (2.5)$$

for some ξ_ρ , BRST-invariant amplitudes are independent of the value of ρ .

Suppose one computes the amplitude $\langle F(\lambda, \bar{\lambda}) \mathcal{N} \rangle$ where $F(\lambda, \bar{\lambda})$ is some BRST-invariant operator. Then ρ -independence implies that the $(-\rho r \theta)^n$ terms in

$$e^{-\rho(\lambda \bar{\lambda} + r \theta)} = e^{-\rho \lambda \bar{\lambda}} \left(1 + \sum_{n=1}^{11} \frac{1}{n!} (-\rho r \theta)^n \right)$$

can only contribute to $\langle F(\lambda, \bar{\lambda}) \mathcal{N} \rangle$ if $\int d^{11} \lambda d^{11} \bar{\lambda} F(\lambda, \bar{\lambda}) e^{-\rho \lambda \bar{\lambda}}$ has poles in ρ . But this implies that $F(\lambda, \bar{\lambda})$ diverges slower than $(\lambda \bar{\lambda})^{-11}$ since

$$\int d^{11} \lambda d^{11} \bar{\lambda} (\lambda \bar{\lambda})^{-n} e^{-\rho \lambda \bar{\lambda}} \sim \rho^{n-11}. \quad (2.6)$$

So θ zero modes in $\mathcal{N}$ can only contribute to $\langle F(\lambda, \bar{\lambda}) \mathcal{N} \rangle$ if $F(\lambda, \bar{\lambda})$ diverges slower than $(\lambda \bar{\lambda})^{-11}$, which implies that $\int d^{11} \lambda d^{11} \bar{\lambda} F(\lambda, \bar{\lambda})$ is well-defined near $\lambda = 0$.

Since the b ghost of (2.2) is spacetime supersymmetric, the sixteen θ^α zero modes in the functional integral of (2.1) must come either from the regulator $\mathcal{N}$ or from the external vertex operators $\prod_{r=1}^N U_r(z_r)$. If all sixteen θ zero modes come from the superfields in the vertex operators U_r , the resulting term in the effective action is not a ten-dimensional F-term since it can be written as an integral over the maximum number of θ 's. However, if at least one of the θ zero modes come from $\mathcal{N}$, the amplitude could contribute to F-terms. Therefore, the above argument implies that amplitudes which contribute to ten-dimensional F-terms do not require regularization when $\lambda \rightarrow 0$.

Note that as in lower dimensions, D=10 F-terms are defined as manifestly gauge-invariant terms in the superspace effective action which cannot be written as integrals over the maximum number of θ 's. Although one does not know how to construct off-shell D=10 superspace actions, one can construct higher-derivative D=10 superspace actions which are functions of on-shell linearized superfields.

For example, for open superstrings, the massless super-Yang-Mills vertex operator is

$$\int dz U = \int dz (A_M(Z) \partial Z^M + W^\alpha(Z) d_\alpha + F_{mn}(Z) N^{mn}) \quad (2.7)$$

where $Z^M = (x^m, \theta^\alpha)$, and the gauge-invariant superfield of lowest dimension is W^α whose lowest component is the gluino of dimension $\frac{1}{2}$. Note that $\int dz U$ can be expressed in manifestly gauge-invariant form by using the normal-coordinate expansion to write

$$\int dz A_M(Z) \partial Z^M = \int dz \left(\frac{1}{2} F_{MN}(Z_0) \hat{Z}^M \partial Z^N + \frac{1}{6} \partial_P F_{MN}(Z_0) \hat{Z}^P \hat{Z}^M \partial Z^N + \dots \right) \quad (2.8)$$

where $\hat{Z}^M = Z^M - Z_0^M$ and Z_0^M is a constant. Since N=1 D=10 superspace contains 16 θ 's, any term in the superspace action involving M superfields W^α which is integrated over the full superspace has dimension $\geq (M + 16)/2$. Therefore, any term in the N=1 D=10 superspace action involving M field-strengths which has dimension less than $(M + 16)/2$ is necessarily an N=1 D=10 F-term.

For closed Type IIB superstrings, the massless supergravity vertex operator is

$$\int d^2z U = \int dz ((G_{MN}(Z) + B_{MN}(Z)) \partial Z^M \bar{\partial} Z^N + W^{\alpha\beta}(Z) d_\alpha \bar{d}_\beta + \dots) \quad (2.9)$$

where $Z^M = (x^m, \theta^\alpha, \bar{\theta}^\beta)$, and the gauge-invariant superfield of lowest dimension is $W^{\alpha\beta}(x, \theta, \bar{\theta})$ whose lowest component is the Ramond-Ramond field strength of dimension 1. Note that the dilaton and axion are gauge-invariant fields of dimension zero, but they always appear with derivatives in the massless vertex operator. Since N=2 D=10 superspace contains 32 θ 's, any term in the superspace action involving M superfields $W^{\alpha\beta}$ which is integrated over the full superspace has dimension $\geq (M + 16)$. Therefore, any term in the N=2 D=10 superspace action involving M field-strengths which has dimension less than $(M + 16)$ is necessarily an N=2 D=10 F-term. For example, since the curvature tensor R_{mnpq} has dimension 2, the term

$$\int d^{10}x \sqrt{g} \partial^L R^M \quad (2.10)$$

in the Type II effective action is an N=2 D=10 F-term if $L + 2M < M + 16$, i.e. if $L + M < 16$.

3. New Multiloop Theorems

The multiloop theorems in this paper will be proven by counting fermionic zero modes in the integrand of (2.1). The left-moving fermionic worldsheet fields in the non-minimal formalism include the superspace variable θ^α and its conjugate momentum d_α , and the non-minimal variable r_α and its conjugate momentum s^α . The non-minimal variable r_α is constrained to satisfy $\bar{\lambda} \gamma^m r = 0$ where $\bar{\lambda} \gamma^m \bar{\lambda} = 0$, so r_α has 11 independent components. Since θ^α and r_α are worldsheet scalars, on a genus g surface θ^α contains 16 zero modes, d_α contains $16g$ zero modes, r_α contains 11 zero modes, and s^α contains $11g$ zero modes. So the amplitude of (2.1) vanishes unless all of these fermionic zero modes are present in the integrand of (2.1).

3.1. Nonrenormalization of $\partial^n R^4$

Using the prescription of (2.1), it will now be proven that perturbative contributions to $\partial^n R^4$ terms vanish above $n/2$ loops. This is proven by showing that the massless four-point g -loop amplitude at low energies is proportional to

$$\left(\frac{\partial}{\partial\theta}\frac{\partial}{\partial\bar\theta}\right)^{2g+4}W^4 = \partial^{2g}R^4 + \dots \quad (3.1)$$

where W is the Ramond-Ramond field strength of (2.9). So $\partial^n R^4$ terms only get perturbative contributions up to genus $n/2$. When $g \geq 6$, (3.1) is no longer an F-term, so there may be $\lambda \rightarrow 0$ divergences which need to be regularized. The theorem has therefore only been proven when $n < 12$.

To get a non-vanishing four-point g -loop amplitude, the integrand of (2.1) must provide $16g$ d_α zero modes which can come either from the four vertex operators of (2.9), from the regulator $\mathcal{N}$ of (2.3), or from the $3g - 3$ b ghosts of (2.2). The most efficient way to obtain these $16g$ zero modes is if the four vertex operators provide the term $(W^{\alpha\beta}d_\alpha\bar{d}_\beta)^4$, and the regulator $\mathcal{N}$ provides the term $(sd)^{11g}$, where $11g$ is the maximum power since there are only $11g$ independent s zero modes.

The remaining $5g - 4$ d_α zero modes must come from the $3g - 3$ b ghosts, and to minimize the number of θ^α zero modes coming from the vertex operators, it will be advantageous to minimize the number of r_α zero modes coming from the b ghosts. The ghost contribution which provides $5g - 4$ d_α zero modes while minimizing the number of r_α zero modes is

$$\left(\frac{\partial x^m(\bar{\lambda}\gamma_m d)}{2(\bar{\lambda}\lambda)}\right)^{g-2} \left(\frac{(\bar{\lambda}\gamma^{mnp}r)(d\gamma_{mnp}d)}{192(\bar{\lambda}\lambda)^2}\right)^{2g-1}, \quad (3.2)$$

where $g - 2$ b ghosts provide the first term and $2g - 1$ b ghosts provide the second term.

The contribution of (3.2) provides $2g - 1$ of the 11 r_α zero modes, so the remaining $12 - 2g$ r_α zero modes must come from $\mathcal{N}$ through the term $(r\theta)^{12-2g}$. Since this term provides $12 - 2g$ of the 16 θ^α zero modes, the remaining $2g + 4$ θ^α zero modes must come from the superfields $W^{\alpha\beta}$ in the vertex operators.

So for the amplitude to be non-vanishing, the four external vertex operators must provide at least $2g + 4$ θ^α zero modes. Therefore, at low energies, the four-point g -loop scattering amplitude is proportional to (3.1) as claimed. Note that it is assumed that there are no inverse factors of momentum coming from the $\langle \prod_{r=1}^4 e^{ik_r x(z_r)} \rangle$ correlation function which would decrease the number of derivatives on R^4 in (3.1). This assumption is reasonable since the massless three-point multiloop amplitude vanishes, so one does not expect any poles in momentum for the four-point multiloop amplitude.

3.2. Equivalence of IIA and IIB $\partial^n R^4$ terms

It will now be proven that up to four loops, four-point graviton contributions to F-terms coincide in the IIA and IIB effective actions. Using the previous theorem that $\partial^n R^4$ terms do not get perturbative contributions above $n/2$ loops, this implies that perturbative contributions to $\partial^n R^4$ terms coincide in the IIA and IIB effective actions for $n \leq 8$.

To prove this multiloop theorem, similar methods to [14] will be used. IIA and IIB superstrings are related by a parity operation on the left-moving worldsheet variables which flips the chirality of the left-moving spacetime spinor. For graviton scattering amplitudes, this parity operation flips the sign of terms which involve an $\epsilon_{m_1 \dots m_{10}}$ tensor coming from the integration over the left-moving variables.

For four-point graviton amplitudes, the only way to contract the vector indices on such an $\epsilon_{m_1 \dots m_{10}}$ tensor is if there is also an $\epsilon_{n_1 \dots n_{10}}$ tensor coming from the integration over the right-moving variables. One can then contract the vector indices of the left-moving ϵ tensor either with the indices of the right-moving ϵ tensor or with the external momenta and polarizations.

Since there are three independent momenta, k_r^m for $r = 1$ to 3, and four independent polarizations, h_r^{mn} for $r = 1$ to 4, the minimum number of indices which must be contracted between the left and right-moving ϵ tensors is three. This can be accomplished using the contraction

$$h_1^{m_1 n_1} h_2^{m_2 n_2} h_3^{m_3 n_3} h_4^{m_4 n_4} k_1^{m_5} k_1^{n_5} k_2^{m_6} k_2^{n_6} k_3^{m_7} k_3^{n_7} \eta^{m_8 n_8} \eta^{m_9 n_9} \eta^{m_{10} n_{10}} \epsilon_{m_1 \dots m_{10}} \epsilon_{n_1 \dots n_{10}}. \quad (3.3)$$

To contract indices of the left and right-moving ϵ tensors, the correlation function must involve factors of ∂x^m and $\bar{\partial} x^n$ since these are the only left and right-moving fields which can be contracted. For example, in the RNS formalism, these factors of ∂x^m and $\bar{\partial} x^n$ come from the left and right-moving picture-changing operators. Since g -loop RNS amplitudes involve $2g - 2$ left and right-moving picture-changing operators, and since one needs at least three ∂x 's and $\bar{\partial} x$'s to perform the contraction of (3.3), the term in (3.3) is only possible when $g \geq 3$. So the four-point graviton amplitudes in IIA and IIB superstring theory have been proven to coincide up to two loops using the RNS formalism [14].

Using the prescription of section 2 for F-term computations, the three factors of ∂x^m and $\bar{\partial} x^n$ can come from the term $\frac{\partial x^m (\bar{\lambda} \gamma_m d)}{2(\bar{\lambda} \lambda)}$ in the b ghost. By counting d_α zero modes as in (3.2), one finds at genus g that the maximum number of ∂x factors is $g - 2$. So

the contraction of (3.3) is only possible when $g \geq 5$, implying that four-point graviton amplitudes contribute equally to IIA and IIB F-terms when $g \leq 4$.

Acknowledgements: I would like to thank Nikita Nekrasov, Jorge Russo, Pierre Vanhove, and especially Michael Green for useful discussions. I would also like to thank the hospitality of Cambridge University and the 2006 Eurostrings conference where a preliminary version of this paper was presented, and CNPq grant 300256/94-9, Pronex grant 66.2002/1998-9, and FAPESP grant 04/11426-0 for partial financial support.

References

- [1] N. Berkovits, *Super-Poincaré covariant quantization of the superstring*, JHEP 0004 (2000) 018, hep-th/0001035.
- [2] N. Berkovits, *Multiloop amplitudes and vanishing theorems using the pure spinor formalism for the superstring*, JHEP 0409 (2004) 047, hep-th/0406055.
- [3] E.J. Martinec, *Nonrenormalization theorems and fermionic string finiteness*, Phys. Lett. B171 (1986) 189;
S. Mandelstam, *The n loop string amplitude: Explicit formulas, finiteness and absence of ambiguities.*, Phys. Lett. B277 (1992) 82.
- [4] M.B. Green and M. Gutperle, *Effects of D instantons*, Nucl. Phys. B498 (1997) 195, hep-th/9701093.
- [5] M.B. Green and P. Vanhove, *D -instantons, strings and M -theory*, Phys. Lett. B408 (1997) 122, hep-th/9704145.
- [6] N. Berkovits, *Super-Poincaré covariant two-loop superstring amplitudes*, JHEP 0601 (2006) 005, hep-th/0503197;
N. Berkovits and C.R. Mafra, *Equivalence of two-loop superstring amplitudes in the pure spinor and RNS formalisms*, Phys. Rev. Lett. 96 (2006) 011602, hep-th/0509234;
C.R. Mafra, *Four-point one-loop amplitude computation in the pure spinor formalism*, JHEP 0601 (2006) 075, hep-th/0512052.
- [7] N. Berkovits, *Pure spinor formalism as an $N=2$ topological string*, JHEP 0510 (2005) 089, hep-th/0509120.
- [8] N. Berkovits and N. Nekrasov, *Multiloop superstring amplitudes from non-minimal pure spinor formalism*, hep-th/0609012.
- [9] N. Berkovits and C.R. Mafra, *Some superstring amplitude computations with the non-minimal pure spinor formalism*, hep-th/0607187.
- [10] N. Nekrasov, *Lectures at the 23rd Jerusalem Winter School in Theoretical Physics, January 2-5, 2006*, <http://www.as.huji.ac.il/schools/phy23/media2.shtml>.
- [11] M.B. Green, J. Russo and P. Vanhove, *to appear*;
M.B. Green, *talk at Strings 2006*.
- [12] M.B. Green and P. Vanhove, *Duality and higher derivative terms in M theory*, JHEP 0601 (2006) 093, hep-th/0510027.
- [13] M.B. Green and P. Vanhove, *The low-energy expansion of the one loop type II superstring amplitude*, Phys. Rev. D61 (2000) 104011, hep-th/9910056.
- [14] M.B. Green, H. Kwon and P. Vanhove, *Two loops in eleven dimensions*, Phys. Rev. D61 (2000) 104010, hep-th/9910055.
- [15] M.B. Green, M. Gutperle and P. Vanhove, *One loop in eleven dimensions*, Phys. Lett. B409 (1997) 177, hep-th/9706175.
- [16] Z. Bern, L. Dixon, D. Dunbar, M. Perelstein and J.S. Rozowsky, *On the relationship between Yang-Mills theory and gravity and its implication for ultraviolet divergences*, Nucl. Phys. B530 (1998) 401, hep-th/9802162.